NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT CHARTBOOK ON PATIENT SAFETY

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Agency for Healthcare Research and Quality
5600 Fishers Lane
Rockville, MD 20857
www.ahrq.gov

AHRQ Publication No. 23-0032
Updates 21-0012
March 202

ACKNOWLEDGMENTS

The National Healthcare Quality and Disparities Report (NHQDR) is the product of collaboration among agencies across the U.S. Department of Health and Human Services (HHS). Many individuals guided and contributed to this effort. Without their magnanimous support, this chartbook would not have been possible. Specifically, we thank:

Authors:

- AHRQ: Barbara Barton, Darryl Gray, and Cecilia Hahn.
- Cormac Corporation: Pam Phojanakong, Morgan Cappa, Naizam Kamookagath.
- Mathematica Inc. Xiaohong (Sharon) Zhao.

Primary AHRQ Staff: Robert Otto Valdez, David Meyers, Erin Grace, Margie Shofer, Karen Chaves, Barbara Barton, Doreen Bonnett, Darryl Gray, Romsai (Tony) Boonyasai, Cecilia Hahn, Lan Liang, Pradip Muhuri, Caren Ginsberg, and Elma Chowdhury.

AHRQ Patient Safety Team Reviewers: Cindy Brach, Emily Chew, Caren Ginsberg, Monika Haugstetter, Lawrence Reid, Margie Shofer, Tselote Tilahun, Andrea Timashenka.

HHS Giho IRQRQ Paa,a

Preventable admissions and readmissions can result from problems with patient safety or

In April 2020, AHRQ published Making Healthcare Safer III, the third compendium of existing and emerging patient safety best practices. In September 2020, AHRQ and the Institute for Healthcare Improvement copublished the National Action Plan to Advance Patient Safety. AHRQ and several organizations committed to patient safety developed this plan. It focuses on culture, leadership, and governance; patient and family engagement; workforce safety; and learning systems—all foundational needs for safe care.

The body of research examining disparities in patient safety continues to evolve in the United States and abroad (Metersky, et al., 2011; Piccardi, et al., 2018; Noursi, et al., 2020; Fasano, et al., 2020; Thomas, et al., 2020).

CHARTBOOK CONTENT

This chartbook includes:

- Summaries of trends across measures of patient safety from the NHQDR.
- Figures illustrating select measures of patient safety.
- Supplemental descriptions and data on patient safety measures from several outside sources.

References:

- Introduction and Methods contains information about methods used in the chartbook.
- A Data Query tool provides access to most NHQDR data tables (https://datatools.ahrq.gov/nhqdr).

Data Sources:

- Agency for Healthcare Research and Quality (AHRQ):
  - Healthcare Cost and Utilization Project (HCUP)
  - Medical Expenditure Panel Survey

5. Postoperative acute kidney injury requiring dialysis per 1,000 elective-surgery admissions, age 18 and over
6. Home health care patients whose surgical wound improved
7. Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age less than 18 years
8. Hospital admissions with iatrogenic pneumothorax per 1,000 medical and surgical admissions, age 18 and over
9. Deaths per 1,000 elective-surgery admissions who developed serious treatable complications of care during hospitalization, ages 18-89 or obstetric admissions
10. Deaths per 1,000 hospital admissions with expected low mortality, age 18 and over or obstetric admissions
11. Home health care patients whose management of oral medications improved
12. High-risk, long-stay nursing home residents with pressure ulcer
13. Long-stay nursing home residents with a urinary tract infection
14. Short-stay nursing home patients with pressure ulcers that are new or worsened

- Not Changing:

1. Perioperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 surgical admissions, age 18 and over
2. Postoperative hip fracture per 1,000 surgical admissions who were not susceptible to falling, age 18 and over
3. Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, age 18 and over
4. Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age 18 and over
5. Birth trauma - injury to neonate per 1,000 live births
6. Long-stay nursing home residents experiencing one or more falls with major injury

Process Measures:

- Improving:

1. Adults age 65 and over who received in the calendar year at least 1 of 11 prescription medications that should be avoided in older adults
2. Adults age 65 and over who received in the calendar year at least 1 of 33 potentially inappropriate prescription medications for older adults
3. Low-risk, long-stay nursing home residents with a catheter inserted and left in the bladder

- Not Changing:

1. Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
2. Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care

3. Adults who reported that home health care providers talked with them about the purpose of taking their new or changed prescription medicines in the last 2 months of care
4. Adults who reported that home health care providers talked with them about when to take medicines in the last 2 months of care
5. Adults who reported that home health care providers talked with them about the side effects of medicines in the last 2 months of care

- Worsening:

1. Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care

Trends by Sub-Area

Number and percentage of patient safety measures improving, not changing, or worsening from 2002 to 2020, by sub-area

- Home Health Communication is the only area in which any measure was worsening:
  - "Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care," which declined from 78.8% in 2012 to 74.5% in 2020.

Complications of Medication:

- Improving:

1. Home health care patients whose management of oral medications improved

Surgical Care:

- Improving:

1. Postoperative pulmonary embolism or deep vein thrombosis per 1,000 surgical admissions, age 18 and over
2. Postoperative respiratory failure, prolonged mechanical ventilation, or reintubation per 1,000 elective-surgery admissions, age 18 and over
3. Postoperative acute kidney injury requiring dialysis per 1,000 elective-surgery

Summary of Patient Safety Trends

Improving

- Measure years are from 2002, 2013, or 2014 through 2019 or 2020.
- Improving measures are defined as average percentage changes greater than 1% per year in a favorable direction and statistically significant.
- The measure of [illegible]

Not Changing

- Measure years are from 2002, 2013, or 2014 through 2019 or 2020.
- Measures not changing are defined as rates of change that are no greater than 1% per year (positive or negative) or are not statistically significant.

Of 29 measures, 11 were not changing. Of the measures not changing over time, five were HHCAHPS measures regarding home health communication about medication:

- Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
- Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care
- Adults who reported that home health care providers talked with them about the purpose of taking their new or changed prescription medicines in the last 2 months of care
- Adults who reported that home health care providers talked with them about when to take medicines in the last 2 months of care
- Adults who reported that home health care providers talked with them about the side effects of medicines in the last 2 months of care

One MDS measure examined nursing home care:

- Long-stay nursing home residents experiencing one or more falls with major injury

The remaining five measures that were not changing were HCUP measures covering surgical care and maternal health:

- Perioperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 surgical admissions, age 18 and over
- Postoperative hip fracture per 1,000 surgical admissions who were not susceptible to falling, age 18 and over
- Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, age 18 and over
- Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age 18 and over
- Birth trauma - injury to neonate per 1,000 live births

Worsening

- Worsening measures are defined as having an AAPC of more than 1% per year in an unfavorable direction and statistically significant.
- Of 29 measures, only one was worsening overall:

DISPARITIES IN PATIENT SAFETY

Overall Disparities

Numb er and percentage of patient safety measures for which members of select groups experienced better, same, or worse quality of care compared with reference group, 2019- 2020

Key: AI/AN = American Indian or Alaska Native; NHPI = Native Hawaiian/Pacific Islander; n = number of measures.
Note: Poor indicates family income less than the federal poverty guideline. High Income indicates family income four times the federal poverty guideline or greater. White, Black, Asian, AI/AN, and NHPI are non-Hispanic. Hispanic includes all races.
Numbers of measures differ across groups because of data limitations. The data shown here are from 2019 or later.
This figure reflects the most current data year available and is not limitTc 0 Tw 43rnlbletionso 76.8 (y)-20.4 3 (o)0. neb ei7.5 ea on0.

3.

9. Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
10. Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care
11. Adults who reported that home health care providers talked with them about the purpose of taking their new or changed uHha[o033 e(n)66.1t([o033 e() c)10.3 (hoM2vi)44.4 1dPu10.34 (ki)4
    -

- Worse Performance:

1. Home health care patients whose surgical wound improved
2.

6. Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care

x

Trends in Disparities

Trends in disparities over time are only examined when a disparity existed in the earliest year of data available. Trends are assessed using unweighted regression. The average annual change (AAC) is computed for the selected group and the reference group. The difference between the AAC for the selected group and the AAC for the reference group is calculated and its statistical

Infections counted for SIRs are restricted to acute care hospitals (excluding critical access hospitals, long-term acute care hospitals, and inpatient rehabilitation facilities) and are stratified by unit type:

- Critical care units (excluding neonatal intensive care units)
- General hospital wards

Data were submitted to NHSN by hospitals in all 50 states, District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands. SIRs were not calculated for states or territories with fewer than five facilities reporting data.

Guam and the Virgin Islands had too few hospitals for the calculation of state-level SIRs for any of the measures presented here. For the same reason, SIRs were not calculated for Vermont in 2017 or 2018 for "Central line-associated bloodstream infections seen in critical care units" and "Catheter-associated urinary tract infections seen in critical care units." Puerto Rico was not included in any year for "Hospital-onset Cloere nareJ / 10.4

Standardized Infection Ratios for Catheter-Associated Urinary Tract Infections

National SIR for catheter-associated urinary tract infections seen in critical care units and wards (non-critical care units), 2015-2020



Source: Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases, National and State Healthcare-Associated Infections Data Report, 2015-2020. https://www.cdc.gov/hai/data/portal/progress-report.html.

Note: SIRs below 1.0 are better. This measure is not included in the summary analysis.

x **Importance:** Compared with rates of other hospital-acquired infections, CAUTI rates vary more among u4.3 (f)-7.8 , (mo7 (e)21.4 (c)4tt)44. 0 Td( )Tj-0.011 Tc 0.011 Tw 0.2 0 Td(in)17.3. Tc 0 Tw 2-.ar

e v 78 , (mo7 (e)21.430.011 Tw 02-.a)-d[(m)-22 im

State-Specific Infection Ratios for CAUTIs in Critical Care Units

State-specific distributions of SIRs for catheter-associated urinary tract infections seen in critical care units, 2015-2020



Source: Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases, National and State Healthcare-Associated Infections Data Report, 2015-2020. http://www.cdc.gov/hai/data/portal/progress-report.html.
Note: SIRs below 1.0 are better. The SIR fo

State-Specific Distribution of Infection Ratios for CAUTIs in Wards

State-specific distributions of SIRs for catheter-associated urinary tract Infections seen on wards (non-critical care units), 2015-2020

Sout ca Sical cawc (e)-1 (r)-3 (s)2 (ca-6 (o)0.7 (ca-2 (s) D8 (i)-9 (r)2 (c.7 (cal)-a2 (s)T4 -7 (cal)- 3 (i)-4 (ca0 (s)t9 (r)-3 (s)-6 (n)l9 (r) a

- Findings: In 2020, the national C. difficile SIR was 0.518 hospitalwide. The 95% confidence intervals for all years follow:
  - 2015, 0.987-0.999
  - 2016, 0.915-0.926
  - 2017, 0.799-0.810
  - 2018, 0.706-0.716
  - 2019, 0.578-0.587
  - 2020, 0.513-0.523

State-Specific Infection Ratios for C. difficile

State-specific distributions of SIRs for hospital-onset Clostridioides difficile infections seen hospitalwide, 2015-2020



- Intended Users: Hospital facilities.
- Available Tools: Checklists, preventable incidence calculators, audit forms, event report templates.
- Potential Measures of Effectiveness:
  - Number of CLABSIs attributable to each unit per month
  - Days since last CLABSI
- Impact: Through use of the CUSP toolkit and CLABSI tools, more than 100 hospital intensive care units (ICUs) in Michigan nearly eliminated CLABSIs. Nationwide, the use of this toolkit helped more than 1,100 hospital ICUs reduce rates of CLABSI by 40% in aggregate. More information is available at https://www.ahrq.gov/workingforquality/priorities-in-action/michigan-health-and-hospital-association-keystone-center.html.

Tools for Reducing Catheter-Associated Urinary Tract Infections in Hospitals

- Purpose: To help hospitals prevent CAUTIs and improve safety culture.
- Methods: Implementing evidence-based practical resources and concepts from the Comprehensive Unit-based Safety Program (CUSP).
- Intended Users: Hospital facilities.
- Available Tools: Guides, checklists, webinars, learning modules, data interpretation guides.
- Link: https://www.ahrq.gov/hai/tools/clabsi-cauti-icu/index.html
- Potential Measures of Effectiveness:
  - Number of symptomatic CAUTIs attributable to each unit per month
  - Days since last CAUTI
- Impact: Use of the CUSP for che a0 C4D <</MCID 106 >>BDC 0 g0 Tc 0 TwoMg77gB.7( )i1.267 0a

- Impact: During the ABATE Infection Trial, a protocol involving CHG use was found to reduce bloodstream infections by more than 30% in adult inpatients who were not in intensive care units and who had specific medical devices.

Maternal Morbidity and Mortality

Maternal mortality, defined as the risk of dying from causes associated with childbirth, is considered a sentinel event in that it is a rare and negative maternal outcome (Adams, et al., 2009). The United States has one of the highest maternal mortality rates compared with other high-income industrialized countries, with 17.4 deaths per 100,000 live births in 2018 (CDC, 2020a). About 700 women die from pregnancy-related complications annually. One-third of pregnancy-related deaths occur 1 week to 1 year after delivery. Three in five pregnancy-related deaths are preventable (CDC, 2019).

Persistent racial and ethnic disparities in maternal mortality have also accompanied the rise in maternal deaths. Black women have a pregnancy-related mortality rate 3 times as high as that of non-Hispanic White women (Petersen, et al., 2019).

Like maternal mortality, severe maternal morbidity, which encompasses unintended outcomes of labor and delivery that result in short-term or long-term health issues, has similarly increased in the United States in recent decades (ACOG, Kilpatrick, & Ecker, 2016).

Many cases of maternal morbidity and mortality are potentially preventable. Factors that contribute to these events have been categorized at the patient, provider, health facility, and system level (Petersen, et al., 2019).

Maternal morbidity and mortality measures include:

- In-hospital deaths per 100,000 delivery hospitalizations.
- Severe maternal morbidity per 1,000 delivery hospitalizations.
- Severe postpartum hemorrha,dvez0 Tw (11.9 (t)267 0 Td.9 (t)267 0 Td.9 (t)267 0 Td.9 (t) (l)4-33.7 (r

Merw222.07e m

In-Hospital Maternal Deaths, by Race/Ethnicity

In-hospital deaths per 100,000 delivery hospitalizations, females ages 12-55, by race/ethnicity, 2019



Key: API = Asian or Pacific Islander.

Source: Agency for Healthcare Research and Quality (AHRQ), Healthcare Cost and Utilization Project, State Inpatient Databases, weighted to provide national estimates; and AHRQ Quality Indicators, v2020.1. More information is available at www.qualityindicators.ahrq.gov/Data/HCUP/Publications.aspx#Publications and Healthcare Cost and Utilization Project (HCUP) publications.

In-Hospital Maternal Deaths, by Age

In-hospital deaths per 100,000 delivery hospitalizations, females ages 12-55, by age, 2019



Source: Agency for Healthcare Research and Quality (AHRQ), Healthcare Cost and Utilization Project, State Inpatient Databases, weighted to provide national estimates; and AHRQ Quality Indicators, v2020.1. More information is in Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the 2022 National Healthcare Quality and Disparities Report (https://www.hcup-us.ahrq.gov/

Severe Maternal Morbidity , by Income

Severe maternal morbidity per 1,000 delivery hospitalizations, females ages 12-55, by median income of patient's ZIP C ode, 2019



Source: Agency for Healthcare Research and Quality (AHRQ), Healthcare Cost and Utilization Project, State Inpatient Databases, weighted to provide national estimates; and AHRQ Quality Indicators, v2020.1. More information is in Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the 2022 National Healthcare Quality and Disparities Report (https://www.hcup-us.ahrq.gov/reports/methods/methods.jsp).

Denominator: Inpatient stays for females ages 12.5 (i)86.3 (c)21shc(:)]TJ 9 Tf-0.038 Tc 05sinTd[(M)25.959.2 (e (t)-67.5 (i)87 (t)-6 ds)-33.6

Severe Maternal Morbidity, by Location of Residence

Severe maternal morbidity per 1,000 delivery hospitalizations, females ages 12- 55, by location of residence, 2019



Cent(i)m 0 0 80 >>BDC 7.3 (ge)]TJ -0.32 -.3D 72 >>BDC q 72 425.6 468 25206 Td [uBDC BT

Preeclampsia or Eclampsia, by Income

Preeclampsia or eclampsia per 1,000 delivery hospitalizations, females ages 12-55, by median income of patient's ZIP Code, 2019



Preeclampsia or Eclampsia, by Payment Source

Preeclampsia or e clampsia per 1,000 delivery hospitalizations, females ages 12- 55, by payment source, 2019



Source: Agency for Healthcare Research and Qu h
Me a

Preeclampsia or Eclampsia, by Location of Residence

Preeclampsia or eclampsia per 1,000 delivery hospitalizations, females ages 12-55, by location of residence, 2019



Source: Agency for Healthcare Research and Quality (AHRQ), Healthcare Cost and Utilization Project, State Inpatient Databases, weighted to provide national estimates; and AHRQ Quality Indicators, v2020.1. More information is in Methods Applying AHRQ Quality Indicators to Healthcare Cost and Utilization Project (HCUP) Data for the 2022 National Healthcare Quality and Disparities Report (https://www.hcup-us.ahrq.gov/reports/methods/methods.jsp).

Venous Thromboembolism or Pulmonary Embolism , by Race/Ethnicity

Venous thromboembolism or pulmonary embolism per 1,000 delivery discharges, females ages 12-55, by race/ethnicity, 2019



Venous Thromboembolism or PulmonarrronarThromonar4Thr(at)22 (ib.3 (haf -0.02 Tw 12 0 0 12 72

Adverse Drug Events

An estimated one-third of all adverse events that occur in the inpatient setting are adverse drug events (ODPHP, 2020a). The HHS National Action Plan for Adverse Drug Event Prevention targets three areas:

- Bleeding related to use of anticoagulants.

related rio.u]TJ 0 Tc 0 Tw 3.133 0 2( )Tj 0.033 Tc -0.033 Tw 0.2 0 Td

- Importance: Hypoglycemic agents ingested by mouth are typically used in patients with type 2 diabetes to control blood sugar levels. In some cases, diabetic patients use hypoglycemic agents together with insulin. The risk of chronic kidney disease increases for people with diabetes, and renal impairment can increase the risk of adverse events related to hypoglycemic agents.
- Groups With Disparities:
  - In 2020,