This document is in the public domain and may be used and reprinted without permission. Citation of the source is appreciated. Suggested citation: National Healthcare Quality and Disparities Report Chartbook on Patient Safety. Rockville, MD: Agency for Healthcare Research and Quality; March 2023. AHRQ Pub. No. 23-0032.

NATIONAL HEALTHCARE QUALITY AND DISPARITIES REPORT
CHARTBOOK ON PATIENT SAFETY

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Agency for Healthcare Research and Quality
5600 Fishers Lane
Rockville, MD 20857
www.ahrq.gov

AHRQ Publication No. 23-0032
Updates 21-0012
March 202

ACKNOWLEDGMENTS

The National Healthcare Quality and Disparities Report (NHQDR) is the product of collaboration among agencies across the U.S. Department of Health and Human Services (HHS). Many individuals guided and contributed to this effort. Without their magnanimous support, this chartbook would not have been possible. Specifically, we thank:

- AHRQ: Barbara Barton, Darryl Gray, and Cecilia Hahn.
- Cormac Corporation: Pam Phojanakong, Morgan Cappa, Naizam Kamookagath.
- Mathematica Inc. Xiaohong (Sharon) Zhao.

Robert Otto Valdez, David Meyers, Erin Grace, Margie Shofer, Karen Chaves, Barbara Barton, Doreen Bonnett, Darryl Gray, Romsai (Tony) Boonyasai, Cecilia Hahn, Lan Liang, Pradip Muhuri, Caren Ginsberg, and Elma Chowdhury.

Cindy Brach, Emily Chew, Caren Ginsberg, Monika Haugstetter, Lawrence Reid, Margie Shofer, Tselote Tilahun, Andrea Timashenka.

INTRODUCTION

This Patient Safety Chartbook is part of a family of documents and tools that support the *National Healthcare Quality and Disparities Report* (NHQDR). The NHQDR is an

Preventable admissions and readmissions can result from problems with patient safety or

In April 2020, AHRQ published Making Healthcare Safer III, the third compendium of existing and emerging patient safety best practices. In September 2020, AHRQ and the Institute for Healthcare Improvement copublished the National Action Plan to Advance Patient Safety. AHRQ and several organizations committed to patient safety developed this plan. It focuses on culture, leadership, and governance; patient and family engagement; workforce safety; and learning systems—all foundational needs for safe care.

The body of research examining disparities in patient safety continues to evolve in the United States and abroad (Metersky, et al., 2011; Piccardi, et al., 2018; Noursi, et al., 2020; Fasano, et al., 2020; Thomas, et al., 2020).

CHARTBOOK CONTENT

This chartbook includes:

- Summaries of trends across measures of patient safety from the NHQDR.
- Figures illustrating select measures of patient safety.
- Supplemental descriptions and data on patient safety measures from several outside sources.

References:

- Introduction and Methods contains information about methods used in the chartbook.
- A Data Query tool provides access to most NHQDR data tables (https://datatools.ahrq.gov/nhqdr).

Data Sources:

- Agency for Healthcare Research and Quality (AHRQ):
  - Healthcare Cost and Utilization Project (HCUP)
  - Medical Expenditure Panel Survey

5. Postoperative acute kidney injury requiring dialysis per 1,000 elective-surgery admissions, age 18 and over
6. Home health care patients whose surgical wound improved
7. Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age less than 18 years
8. Hospital admissions with iatrogenic pneumothorax per 1,000 medical and surgical admissions, age 18 and over
9. Deaths per 1,000 elective-surgery admissions who developed serious treatable complications of care during hospitalization, ages 18-89 or obstetric admissions
10. Deaths per 1,000 hospital admissions with expected low mortality, age 18 and over or obstetric admissions
11. Home health care patients whose management of oral medications improved
12. High-risk, long-stay nursing home residents with pressure ulcer
13. Long-stay nursing home residents with a urinary tract infection
14. Short-stay nursing home patients with pressure ulcers that are new or worsened

- Not Changing:

1. Perioperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 surgical admissions, age 18 and over
2. Postoperative hip fracture per 1,000 surgical admissions who were not susceptible to falling, age 18 and over
3. Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, age 18 and over
4. Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age 18 and over
5. Birth trauma - injury to neonate per 1,000 live births
6. Long-stay nursing home residents experiencing one or more falls with major injury

- Improving:

1. Adults age 65 and over who received in the calendar year at least 1 of 11 prescription medications that should be avoided in older adults
2. Adults age 65 and over who received in the calendar year at least 1 of 33 potentially inappropriate prescription medications for older adults
3. Low-risk, long-stay nursing home residents with a catheter inserted and left in the bladder

- Not Changing:

1. Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
2. Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care

3. Adults who reported that home health care providers talked with them about the purpose of taking their new or changed prescription medicines in the last 2 months of care
4. Adults who reported that home health care providers talked with them about when to take medicines in the last 2 months of care
5. Adults who reported that home health care providers talked with them about the side effects of medicines in the last 2 months of care

- Worsening:

1. Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care

Trends by Sub-Area

Number and percentage of patient safety measures improving, not changing, or worsening from 2002 to 2020, by sub-area

- Home Health Communication is the only area in which any measure was worsening:
  - "Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care," which declined from 78.8% in 2012 to 74.5% in 2020.

- Improving:

  1. Home health care patients whose management of oral medications improved

- Improving:

  1. Postoperative pulmonary embolism or deep vein thrombosis per 1,000 surgical admissions, age 18 and over
  2. Postoperative respiratory failure, prolonged mechanical ventilation, or reintubation per 1,000 elective-surgery admissions, age 18 and over
  3. Postoperative acute kidney injury requiring dialysis per 1,000 elective-surgery

Summary of Patient Safety Trends

Improving

- Measure years are from 2002, 2013, or 2014 through 2019 or 2020.
- Improving measures are defined as average percentage changes greater than 1% per year in a favorable direction and statistically significant.
- The measure of [illegible]

Not Changing

- Measure years are from 2002, 2013, or 2014 through 2019 or 2020.
- Measures not changing are defined as rates of change that are no greater than 1% per year (positive or negative) or are not statistically significant.

Of 29 measures, 11 were not changing. Of the measures not changing over time, five were HHCAHPS measures regarding home health communication about medication:

- Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
- Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care
- Adults who reported that home health care providers talked with them about the purpose of taking their new or changed prescription medicines in the last 2 months of care
- Adults who reported that home health care providers talked with them about when to take medicines in the last 2 months of care
- Adults who reported that home health care providers talked with them about the side effects of medicines in the last 2 months of care

One MDS measure examined nursing home care:

- Long-stay nursing home residents experiencing one or more falls with major injury

The remaining five measures that were not changing were HCUP measures covering surgical care and maternal health:

- Perioperative hemorrhage or hematoma with surgical drainage or evacuation per 1,000 surgical admissions, age 18 and over
- Postoperative hip fracture per 1,000 surgical admissions who were not susceptible to falling, age 18 and over
- Reclosure of postoperative abdominal wound dehiscence per 1,000 abdominopelvic-surgery admissions of length 2 or more days, age 18 and over
- Accidental puncture or laceration during a procedure per 1,000 medical and surgical admissions, age 18 and over
- Birth trauma - injury to neonate per 1,000 live births

Worsening

- Worsening measures are defined as having an AAPC of more than 1% per year in an unfavorable direction and statistically significant.
- Of 29 measures, only one was worsening overall:

DISPARITIES IN PATIENT SAFETY

Overall Disparities

Number and percentage of patient safety measures for which members of select groups experienced better, same, or worse quality of care compared with reference group, 2019-2020

AI/AN = American Indian or Alaska Native; NHPI = Native Hawaiian/Pacific Islander; n = number of measures.

Poor indicates family income less than the federal poverty guideline. High Income indicates family income four times the federal poverty guideline or greater. White, Black, Asian, AI/AN, and NHPI are non-Hispanic. Hispanic includes all races.

Numbers of measures differ across groups because of data limitations. The data shown here are from 2019 or later.

This figure reflects the most current data year available and is not limit Tc 0 Tw 43rnlbletionso 76.8 (y)-20.4 3 (o)0. neb ei7.5 ea on0.

3.

9. Adults who reported a home health care provider talking with them about how to set up their home so they can move around safely when they first started getting home health care
10. Adults who reported a home health care provider talking with them about all the prescription and over-the-counter medicines they were taking when they first started getting home health care
11. Adults who reported that home health care providers talked with them about the purpose of taking their new or changed uHha[o033 e(n)66.1t([o033 e() c)10.3 (hoM2vi)44.4 1dPu10.34 (ki)4
    -

- Worse Performance:

    1. Home health care patients whose surgical wound improved
    2.

6. Adults who reported a home health care provider asking to see all the prescription and over-the-counter medicines they were taking when they first started getting home health care

Trends in Disparities

Trends in disparities over time are only examined when a disparity existed in the earliest year of data available. Trends are assessed using unweighted regression. The average annual change (AAC) is computed for the selected group and the reference group. The difference between the AAC for the selected group and the AAC for the reference group is calculated and its statistical