

For a medical

Search Description	Objective	Subcategories	Citations
		SR/MA	1

Liquid	

RCT	Randomized controlled trial
RNA	Ribonucleic acid
SR	Systematic review

Key Informantsc 0.002 Tw -11c 661.68 198.36 065.26c reW16w 2.92vt0.88 sAMCID 26- BDC - 0.06

Name	Theo deVos
Affiliation	Epigenomics Inc.
Key Question 1	KQ1: Currently, KQ1 refers to “adults at risk for lung, prostate, breast, ovarian or colorectal cancer.”

Name	Trisha Brown
Affiliation	Illumina

	The scope of clinical use for liquid biopsy is large. While the background accurately describes the potential for screening, diagnosis and management of cancer, the majority of commercially available tests and clinical evidence published to date are focused on the management of cancer using cfDNA. To make this review relevant, timely and streamlined in terms of appropriate comparators, clinical validity, and clinical utility, it is recommended to focus solely on the management of cancer using cell free DNA, and consider screening and diagnosis, and circulating tumor cells, in separate technology assessments in the future. We believe that including screening and diagnosis applications and CTC would be premature at this point in time. Focusing the technology assessment will make the final report more consumable by end users as well. In addition, there are a significant number of tests commercially available and in the pipeline for pan-cancer solid tumor applications, as well as individual cancers such as bladder, renal, and heme. These are important considerations for the Medicare population. Therefore, it is recommended that a pan-cancer approach, more narrowly focused on cancer management and cfDNA, be considered for this technology assessment.
Title	
Duration	

Title	Sincerely, Brian Alexander, MD, MPH Chief Medical Officer Foundation Medicine, Inc.

Name	Miguel R. Ossandon
Affiliation	NIH/NCI/Division of Cancer Treatment and Diagnosis/Cancer Diagnosis Program
Key Question 1	
Key Question 2	
Key Question 3	
Analytic Framework	On figure 1B is reed: "Adult at high risk or suspected to have lung, prostate, breast, ovarian, or colorectal cancer". Comment: It is not necessary to include "high risk" in the diagnosis paradigm. If the patient is

Name	
Affiliation	Guardant Health
Key Question 1	
Key Question 2	
Key Question 3	
Analytic Framework	The proposed analytic framework does not take into account rapidly evolving body of research on liquid biopsy. A thorough examination of the literature must acknowledge research conducted outside of the U.S. and reflect the significant advances made since the previous topic refinement in October 2018. As liquid biopsy becomes more widely used, any evaluation

	(Mok T, Wu YL, Lee JS, et al. Detection and Dynamic Changes of EGFR Mutations from Circulating Tumor DNA as a Predictor of Survival Outcomes in NSCLC Patients Treated with First-line Intercalated Erlotinib and Chemotherapy. Clin Cancer Res 2015;21:3196-203.) (Nakamura Y. Yoshino T. Clinical Utility of Analyzing Circulating Tumor DNA in Patients With Metastatic Colorectal Cancer. Oncologist. 2018 Nov;23(11):1310-1318. doi: 10.1634/theoncologist.2017-0621. Epub 2018 Apr 26.) In 2018, Aggarwal et al. at the University of Pennsylvania conducted a retrospective analysis of the real-world use of Guardant360 in the care of 323 consecutive metastatic non-small cell lung cancer patients, half of which were tested at the first line. First, the study found that tissue genotyping failed in nearly half (44%) of the 71% of patients in whom the physician elected to use tissue genotyping, which demonstrates how large of a challenge tissue-based genotyping presents in a real-world setting. Secondly, in those patients for whom both tissue and plasma results were available at diagnosis, concordance of the two was very high, at nearly 90%. And thirdly, the addition of Guardant360 nearly doubled the proportion of patients identified with standard of care actionable biomarkers from 21% to 36%, further demonstrating that real-world use of Guardant360 substantially increases the number of treatment opportunities for patients. Finally, and perhaps most importantly, 86% of patients treated based on Guardant360 results achieved either a complete response, partial response or stable disease further demonstrating the high clinical accuracy of Guardant360 testing. These results, combined with patient satisfaction with the relative ease of providing blood rather than a solid tissue sample, suggest a clinical strategy of pursuing plasma NGS first, then tissue NGS if plasma NGS cannot detect relevant mutations. (Aggarwal C et al. Clinical Implications of Plasma-Based Genotyping With the Delivery of Personalized Therapy in Metastatic Non-Small Cell Lung Cancer. JAMA Oncology 2019 Feb 1;5(2):173-180. doi: 10.1001/jamaoncol.2018.4305.) There are inherent challenges in using tissue biopsy as the reference standard for liquid biopsy. Many patients have no access to receive tumor biopsy due to the invasiveness of the procedure or the tumor tissue obtained is not sufficient for detection of gene alterations. Moreover, tumor heterogeneity makes the tumor biopsy in one site not able to cover the comprehensive genomic profiles. (Wu, Z et al Update on liquid biopsy in clinical management of non-small cell lung cancer. OncoTargets and Therapy 2019:12, 5098) (Sai-Hong Ignatius Ou. Liquid Biopsy to Identify Actionable Genomic Alterations. American Society of Clinical Oncology Educational Book 38 (May 23, 2018) 978-997. DOI: 10,1200/EDBK_199765.) In many regards, liquid biopsy is proven to produce results that tissue cannot. In evaluating the role and clinical utility of liquid biopsy, we maintain that AHRQ will need to determine the correct comparator - and the most appropriate modeling or statistical technique to apply to ensure that the relative benefits of each test are able to be compared indirectly, the example of plain radiography compared to computed tomography (CT) imaging is a relevant example. Tissue biopsy alone is insufficient. What process has AHRQ used for other emerging technologies to avoid comparing to an outdated standard?
Outcomes	When compared to tissue, liquid biopsy has a significantly shorter turnaround time. A 2018 prospective, multicenter, head-to-head study of SOC tissue-based genomic testing to plasma-based comprehensive cfDNA genomic testing revealed liquid biopsy results via Guardant360 were returned significantly faster than tissue, by almost a week. The median turnaround time

patient's medical history, physical examination, and laboratory test results. We therefore request AHRQ to clarify whether "diagnosis" is the correct intended use for KQ2.

A more appropriate application of liquid biopsies may be for surveillance in high-risk populations, but only where such surveillance already has demonstrated clinical utility. As with screening in KQ1, the patient management pathways and clinical utility of surveillance differ for different cancers. For example, the clinical utility of surveillance for CRC in patients diagnosed with hereditary nonpolyposis colorectal cancer (HNPCC) or familial adenomatous

	KQ3 (e.g., MRD or recurrence testing) will vary for different types of cancer, even among the five cancers listed. Reference(s): 1. https://www.fda.gov/medical-devices/vitro-diagnostics/list-cleared-or-approved-companion-diagnostic-devices-vitro-and-imaging-tools
Analytic Framework	As stated above, there appears to be a discrepancy between the language in KQ2, which describes the intended use population as "adults suspected to have lung, prostate, breast, ovarian, or colorectal cancer" and Figure 1b, which describes the same population as "adults at high risk or suspected to have lung, prostate, breast, ovarian, or colorectal cancer." These J0 Tc 0 T (e)1.i should be reconciled for c

	describes a test whose intended use is unclear because the patient population for this case-control study is unclear. 9. In addition to the challenge of intra-tumoral heterogeneity, clonal hematopoiesis of indeterminate potential (CHIP) should also be addressed and represents a source of somatic mutations that limit the specificity of ctDNA mutation-based tests. Reference(s): Chabon, J.J., Hamilton, E.G., Kurtz, D.M. et al. Integrating genomic features for non-invasive early lung cancer detection. Nature 580, 245-251 (2020). https://doi.org/10.1038/s41586-020-2140-0
Population(s)	Please see "General Comments," as well as specific comments for KQ1 (comments 1 and 4), KQ1b, KQ2, KQ2b, KQ3, and KQ3b.
Intervention	Please see "General Comments" and Background (comment 2), as well as specific comments for KQ1 (comment 2), KQ1a, KQ2a, and KQ3a
Comparators	Please see "General Comments."
Outcomes	None
Timing	None
Settings	None
Subgroup Analyses	None
References	Reference #4 seems to be an abstract that describes a test whose intended use is unclear because the patient population for this case-control study is unclear.
Comments	On behalf of Freenome, we appreciate the opportunity to submit comments regarding the Key Question Posting Document. At Freenome, our mission is to develop tools to empower everyone to prevent, detect, and treat disease, with an initial focus on colorectal cancer (CRC). Freenome plans to seek FDA approval for a novel artificial intelligence (AI)/machine learning (ML)-enabled multiomics approach to CRC screening that we believe will allow us to achieve greater clinical sensitivity and specificity than conventional blood-based biomarker approaches, particularly for early-

	screening technologies, such as newer stool-based tests, CT colonography, or colon capsule for CRC; 3D breast tomosynthesis or positron emission mammography for breast cancer; the prostate health index (PHI) or 4Kscore for prostate cancer; and so forth. A thorough evidence review of the analytical validity (AV), clinical validity (CV), and clinical utility(CU) of existing screening strategies is an appropriate and necessary precursor to such an analysis focused on a novel and rapidly evolving science (i.e., liquid biopsies), if only to properly contextualize the analysis. Freenome therefore suggests that AHRQ include such an analysis if KQ1 is retained. Second, the scope of the existing proposal is quite significant, potentially representing every intended use for liquid biopsies in cancer. At a minimum, we encourage AHRQ to focus the proposal on the intended uses for which the peer-reviewed published literature is most mature (i.e., therapy selection and, to a lesser extent, therapeutic monitoring), with an understanding that all uses of liquid biopsies are still relatively nascent, and to contextualize even these analyses with a robust assessment of the AV, CV, and CU of alternative approaches, both standard-of-care and emerging. For example, we are not aware whether similar TAs have been performed for tissue-based testing for therapy selection in late-stage cancer patients, or for existing methods for MRD monitoring or recurrence detection. Finally, one size does not fit all in cancer, whether in the context of screening, therapy selection, or therapeutic monitoring. There are over 100 different types of cancer. The evidentiary requirements for AV, CV, and CU for any test, liquid biopsy or otherwise, necessarily and appropriately differ for different cancers because the risks and benefits of subsequent treatments and interventions differ, and therefore so too do the risks and benefits of screening, surveillance, therapy selection, and therapeutic response monitoring. We therefore urge AHRQ, when performing this evidence review, to consider that the evidentiary requirements for AV, CV, and CU necessarily and appropriately vary for different intended uses in different cancers. Thank you for considering our comments. We look forward to the draft Technology Assessment. Please contact me at XXX should you have any questions or seek additional information. Sincerely yours, Girish Putcha, M.D., Ph.D. Chief Medical Officer Clinical Laboratory Director Freenome
Title	
Duration	

Name	Christopher Keir, MD, MS
Affiliation	GRAIL Inc.
Key Question 1	GRAIL believes that based on evolving technological developments and ongoing clinical studies of liquid biopsy for cancer screening that have not yet been published, it would be premature at this time to include this question and KQ2 within the scope of the Technology Assessment. A comprehensive assessment of a field of science to analyze such metrics as pre-analytic, analytic, clinical validity, and clinical utility to help inform important governmental and regulatory decision-making should be conducted at a time when sufficient scientific information and clinical data are available. GRAIL believes that critical data generation to address and guide the Draft Key Questions is still ongoing. In addition, data supporting the measures of clinical utility of a cancer screening application are distinct and not easily combined with a post-cancer diagnosis application.
Key Question 2	See comments to Question 1.
Key Question 3	
Analytic Framework	
Background	
Population(s)	GRAIL believes it is premature to include liquid biopsy for cancer screening within the scope of the Technology Assessment, and urges AHRQ to remove “patients at risk” and “patients suspected to have” from the analysis population.
Intervention	Circulating tumor cells and circulating tumor DNA are purified from blood and analyzed using

t)08 8 ref15-0.001 Tc 10.98 187.32 43[illegible] (Pericles, the presence of Type I stage 3) 0.5 0.5 0.4 0.2 (4) 2.6 [illegible]

Hamakawa T; “Monitoring Gastric Cancer Progression With Circulating Tumour DNA.” British Journal of Cancer, U.S. National Library of Medicine
Jenkins S; “Plasma CtDNA Analysis for Detection of the EGFR T790M Mutation in Patients With Advanced Non-Small Cell Lung Cancer.” Journal of Thoracic Oncology : Official Publication of the International Association for the Study of Lung Cancer, U.S. National Library of Medicine
Lipson EJ; “Circulating Tumor DNA Analysis as a Real-Time Method for Monitoring Tumor Burden in Melanoma Patients Undergoing Treatment With Immune Checkpoint Blockade.” Journal for Immunotherapy of Cancer, U.S. National Library of Medicine,
M;, Rossi G; “Promises and Pitfalls of Using Liquid Biopsy for Precision Medicine.” Cancer Research, U.S. National Library of Medicine,
Merker JD; “Circulating Tumor DNA Analysis in Patients With Cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review.” Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology, U.S. National Library of Medicine,
Mok T; “Detection and Dynamic Changes of EGFR Mutations From Circulating Tumor DNA as a Predictor of Survival Outcomes in NSCLC Patients Treated With First-Line Intercalated Erlotinib and Chemotherapy.” Clinical Cancer Research : an Official Journal of the American Association for Cancer Research, U.S. National Library of Medicine,
Reinert T; “Analysis of Circulating Tumour DNA to Monitor Disease Burden Following Colorectal Cancer Surgery.” Gut, U.S. National Library of Medicine,
Schmiegel W; “Blood-Based Detection of RAS Mutations to Guide Anti-EGFR Therapy in

different intervention than a test to detect a cancer signal in a normal or high risk individual without known cancer. Moreover, this scope of work is considering a broad range of different

New blood-based molecular diagnostic technologies are emerging that have the potential to address challenges in early detection of cancer (e.g., lack of guideline-recommended screening tests for many types of cancer). These assays, which may allow physicians to detect cancer when outcomes are better and costs are lower, are increasingly supported by foundational evidence establishing their performance. Indeed, member companies regularly present data regarding the performance of these assays at professional society meetings (e.g., American Society for Clinical Oncology (ASCO) and American Association for Cancer Research (AACR)), and this research is increasingly published in peer-reviewed journals. C21 expects there to be a substantial increase in the amount and quality of this data -

For instance, tissue biopsy is not generally used as a disease screening tool. But blood sampling may prove to have utility for the purpose of identifying individuals at average or increased risk for a condition. Similarly, tissue biopsy is not a tool for quantifying the extent of disease; miminal/molecular residual disease assessment and quantification, as well as responsiveness to intervention, can be assessed by blood based sampling and the measurement of circulating tumor DNA (ctDNA). Blood is an ideal sample to detect both germline and somatic DNA variations that can be informative about disease origin, diagnosis, prognosis or susceptibility to treatment and is far simpler, faster and less expensive at getting this information than older tissue biopsy approaches.

For some use cases, the proper comparator for the use of blood (liquid biopsy) as an disease assessment aid is the traditional information gleaned from proper tissue biopsy analysis. In other situations, tissue is not the "gold standard". For instance, in the assessment of MRD using ctDNA, the proper comparator is clinical assessment with imaging supplementation. Both the sensitivity of detection and quantification of tumor burden have been improved with the newer "liquid biopsy" ctDNA based method.